

APPENDIX IV KEY NATIVE ECOSYSTEM PESTS – REGIONAL PEST MANAGEMENT STRATEGY

Argentine ant	<i>(Linepithema humile)</i>
Brown bullhead catfish	<i>(Ameiurensis nebulosis)</i>
European hedgehog	<i>(Erinaceus europaeus occidentalis)</i>
Feral cat	<i>(Felis catus)</i>
Feral goat	<i>(Capra hircus)</i>
Feral pig	<i>(Sus scrofa)</i>
Ferret	<i>(Mustela furo)</i>
Goldfish	<i>(Carassius auratus)</i>
Hare	<i>(Lepus europaeus occidentalis)</i>
House mouse	<i>(Mus musculus)</i>
Koi carp	<i>(Cyprinus carpio)</i>
Mosquito fish	<i>(Gambusia affinis)</i>
Norway rat	<i>(Rattus norvegicus)</i>
Possum	<i>(Trichosurus vulpecula)</i>
Rudd	<i>(Scardinius erythrophthalmus)</i>
Ship rat	<i>(Rattus rattus)</i>
Stoat	<i>(Mustela erminea)</i>
Sulphur crested cockatoo	<i>(Cacatua galerita)</i>
Tench	<i>(Tinca tinca)</i>
Wasp	<i>(Vulpecula germanica; Vulpecula vulgaris)</i>
Weasel	<i>(Mustela nivalis)</i>
African club moss	<i>(Selaginella kraussiana)</i>
Artillery plant	<i>(Galeobdolon luteum)</i>
Barberry	<i>(Berberis glaucocarpa)</i>
Blackberry	<i>(Rubus fruticosus)</i>
Blue morning glory	<i>(Ipomoea indica)</i>
Boxthorn	<i>(Lycium ferocissimum)</i>
Broom	<i>(Cytisus scoparius)</i>
Brush wattle	<i>(Paraserianthes lophantha)</i>
Buddleia	<i>(Buddleja davidii)</i>
Cape honey flower	<i>(Melianthus major)</i>
Cape ivy	<i>(Senecio angulatus)</i>
Chilean flame creeper	<i>(Tropaeolum speciosum)</i>
Chinese and tree privet	<i>(Ligustrum sinense; L. lucidum)</i>
Climbing dock	<i>(Rumex sagittatus)</i>
Great bindweed	<i>(Calystegia silvatica)</i>
Cotoneaster	<i>(Cotoneaster franchetii, C. horizontalis)</i>
Darwin's barberry	<i>(Berberis darwinii)</i>
Egeria	<i>(Egeria densa)</i>
Elaeagnus	<i>(Elaeagnus x reflexa)</i>
German ivy	<i>(Senecio mikanioides)</i>
Gunnera	<i>(Gunnera tinctoria)</i>
Hawthorn	<i>(Crataegus monogyna)</i>

Himalayan honeysuckle	<i>(Leycesteria formosa)</i>
Japanese honeysuckle	<i>(Lonicera japonica)</i>
Lagarosiphon	<i>(Lagarosiphon major)</i>
Marram grass	<i>(Ammophila arenaria)</i>
Mexican daisy	<i>(Erigeron karvinskianus)</i>
Mile-a-minute	<i>(Dipogon lignosus)</i>
Pampas grass	<i>(Cortaderia jubata, C. selloana)</i>
Parrot's feather	<i>(Myriophyllum aquaticum)</i>
Periwinkle	<i>(Vinca major)</i>
Plectranthus	<i>(Plectranthus ciliatus)</i>
Purple ragwort	<i>(Senecio glastifolius)</i>
Spanish heath	<i>(Erica lusitanica)</i>
Stinking iris	<i>(Iris foetidissima)</i>
Sycamore	<i>(Acer pseudoplatanus)</i>
Velvet groundsel	<i>(Senecio petasitis)</i>
Wandering jew	<i>(Tradescantia fluminensis)</i>
Wild onion	<i>(Allium triquetrum)</i>
Wilding conifers	<i>(Larix decidua; var)</i>
Wilding pines	<i>(Pinus spp.)</i>